

**Amendments to the Specification:**

Please replace paragraph [00191] with the following amended paragraph:

[00191] Animal model: The formalin test is a suitable model for the field of inflammatory/persistent pain (~~Dubuisson~~ Dubuisson et al., ~~1997~~ 1977). In this test, 50 µl 5% formalin solution were injected unilaterally into the hind paw of adult Wistar rats and the animals were sacrificed 24 hours after the injection for removal of tissue. In parallel, isotonic saline solution was injected into the hind paw in the control animals.

Please replace paragraph [00228] with the following amended paragraph:

[0001] The following documents are incorporated herein by reference in their entirety.

Akopian AN, Sivilotti L & Wood JN (1995) Nature 379:257-262.

Ausubel FM, Brent R, Kingdon RE, Moore DD, Seidman JG, Smith JA & Struhl K eds. (1990) Current protocols in molecular biology. John Wiley & Sons, Inc. New York, NY.

Baba H, Doubell TP, Woolf CJ 1999: Peripheral inflammation facilitates A $\beta$  fiber-mediated synaptic input to the substantia gelatinosa of the adult rat spinal cord. J Neurosci 19:859-867.

Bauer D, Müller H, Reich J, Riedel H, Ahrenkiel V, Warthoe P & Strauss M (1993): Identification of differentially expressed mRNA species by an improved display technique (DDRT-PCR) Nucl Acids Res 21:4272-4280.

Bonini A, Anderson SM, Steiner DF (1997) Molecular cloning and tissue expression of a novel orphan G protein-coupled receptor from rat lung. *Biochem Biophys Res Comm* 234:190-193.

Chih-Cheng et al., (1995): A P2X prinoceptor expressed by a subset of sensory neurons. *Nature* 377:428-432.

Corderre TJ, Katz J, Vaccarino AL, Melzack R (1993): Contribution of central plasticity to pathological pain: review of clinical and experimental evidence. *Pain* 52:259-285.

Dickenson (1995) Novel pharmacological targets in the treatment of pain. *Pain Rev.*, 2, 1-12.

Dubuisson et al., [1997] 1977 *Pain*, 4:161-174.

Feng Y & Gregor P (1997) Cloning of a novel member of the G protein-coupled receptor family related to peptide receptors. *Biochem Biophys Res Comm* 231:651-654.

Furukawa T, Yang Y, Nakamoto B, Stamatoyannopoulos G, Papayannopoulou T (1966): Identification of new genes expressed in a human erythroleukemia cell line. *Bloods Cell Mol & Dis* 22:11-22.

Gunasekar PG, Kanthasamy, AG, Borowitz JL, Isom GE 1995: NMDA receptor activation produces concurrent generation of nitric oxide and reactive oxygen species: implication for cell death. *J Neurochem* 65:2016-2021.

Hawes BE, Fried S, Yao X, Weig B, Graziano MP 1998: Nociceptin (ORL1) and  $\mu$ -opioid receptors mediate mitogen-activated protein kinase activation in CHO cells through a Gi-coupled signaling pathway: evidence for distinct mechanisms of agonist-mediated desensitization. *J Neurochem* 71:1024-1033.

Hubank M & Schatz DG (1994): Identifying differences in mRNA expression by representational difference analysis of cDNA. *Nucl Acids Res* 22:5640-5648.

- Klußmann S et al., 1996: Nature Biotechnology 14:1112-1115.
- Li L-Y & Chang K-J 1996: The stimulatory effect of opioids on mitogen-activated protein kinase in chinese hamster ovary cells transfected to express  $\mu$ -opioid receptors. Mol Pharm 50:599-602.
- Lian P & Pardee AB 1992: Differential display of eukaryotic messenger RNA by means of the polymerase chain reaction. Science 257:967-971.
- Methner A, Hermey G, Schinke B, Hermanns-Borgmeyer I (1997): A novel G protein-coupled receptor with homology to neuropeptide and chemoattractant receptors expressed during bone development. Biochem Biophys Res Comm 233:336-342.
- Mohit AA, Martin JH & Miller CA 1995: p493F12 kinase: a novel MAP kinase expressed in a subset of neurons in the human nervous system. Neuron 14:67-78.
- Poirier GM-C, Pyati J, Wan JS, Erlander MG 1997: Screening differentially expressed cDNA clones obtained by differential display using amplified RNA. Nucleic Acids Research 25:913-914.
- Sambrook J, Fritsch EF & Maniatis T 1989: Molecular Cloning: A Laboratory Manual. Second Edition. Cold Spring Harbor Laboratory Press, Cold Spring Harbor.
- Sompayrac L, Jane S, Burn TC, Tenen DG & Danna KJ 1995: Overcoming limitations of the mRNA differential display technique. Nucleic Acids Research 23:4738-4739.
- Tal M 1996: A novel antioxidant alleviates heat hyperalgesia in rats with an experimental painful neuropathy. Neurreport 7:1382-1384.
- Tölle TR (1997): Chronischer Schmerz. In: Klinische Neurobiologie [Chronic Pain. In: Clinical Neurobiology], Herdergen T. Tölle TR, Bähr M (eds.): p. 307-336; Spektrum Verlag, Heidelberg.

U.S. Patent No. 5,262,311, issued November 16, 1993.

Velculescu VE, Zhang L, Vogelstein B, Kinzler KW (1995): Serial analysis of gene expression. *Science* 270:484-487.

Wan JS, Sharp JS et al. (1996): Cloning differentially expressed mRNAs. *Nature Biotech* 14:1685-1691.

Watson JB & Margulies JE (1993) Differential cDNA screening strategies to identify novel stage-specific proteins in the developing mammalian brain. *Dev Neurosci* 15:77-86.

Wilks AF (1989) Two putative protein-tyrosine kinases identified by application of the polymerase chain reaction. *Proc Natl Acad Sci USA* 86:1603-1607.

Woolf CJ, Shortland P, Coggeshall RE 1992: Peripheral nerve injury triggers central sprouting of myelinated afferents. *Nature* 355:75-78.

Zimmermann, M & Herdegen, T (1996): Plasticity of the nervous system at the systemic, cellular and molecular levels: a mechanism of chronic pain and hyperalgesia. *Progr Brain Res* 110:233-259.